


















<p>Active ingredients (generic names)</p>	<p>METOCLOPRAMIDE (me-to-clo-pra-mide) PROCHLORPERAZINE (pro-chlor-per-a-zeen)</p> <p></p> <p>Other dopamine antagonists Not included in this monograph — see <i>AMH</i> for information:</p> <ul style="list-style-type: none">• Domperidone• Droperidol• Haloperidol
<p>What it is used for</p>	<ul style="list-style-type: none">• Increasing gastrointestinal movement after operations• Nausea and vomiting (adult) <p>Metoclopramide</p> <ul style="list-style-type: none">• Nausea and vomiting (adult) with: opioid withdrawal, palliative care, pregnancy <p>Prochlorperazine</p> <ul style="list-style-type: none">• Dizziness and vertigo
<p>How it works</p>	<ul style="list-style-type: none">• Acts on dopamine hormone receptors in brain and gut to reduce nausea and vomiting and increase gastrointestinal movement <p></p>
<p>Side effects</p>	<ul style="list-style-type: none">• Restlessness• Agitation• Muscle spasm• Oculogyric crisis <p> Sleepy</p> <p> Constipation</p> <p> Headache (metoclopramide)</p> <p> Dizzy (metoclopramide)</p>

 <p>Warnings</p>	<ul style="list-style-type: none"> • People with Parkinson's disease <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Children</p> </div> <div style="text-align: center;">  <p>Old people</p> </div> <div style="text-align: center;">  <p>Kidney trouble (metoclopramide)</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Dehydration — may need oral rehydration salts</p> </div> <div style="text-align: center;">  <p>Depression</p> </div> <div style="text-align: center;">  <p>Medical consult if needed for more than a short time (one week)</p> </div> </div>
<p>Tell the patient</p> 	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Do not drive or operate machinery if sleepy</p> </div> <div style="text-align: center;">  <p>Drink lots of water</p> </div> <div style="text-align: center;">  <p>Return to clinic if still sick in 1–2 days</p> </div> </div>
<p>Check</p>	<p>Warning stickers (page 310): Metoclopramide 12 Prochlorperazine 1, 13, 16</p> <ul style="list-style-type: none"> • Does patient have signs of dehydration • Check for cause of vomiting