








<p>Active ingredients (generic names)</p>	<p>ONDANSETRON (on-dan-set-ron)</p>  <p>Other 5HT3 antagonists Not included in this monograph — see <i>AMH</i> for information:</p> <ul style="list-style-type: none"> • Granisetron • Palonosetron • Tropisetron
<p>What it is used for</p>	<ul style="list-style-type: none"> • Nausea and vomiting • Persistent vomiting with diarrhoea (children)
<p>How it works</p>	<ul style="list-style-type: none"> • Stops nausea and vomiting by blocking central and peripheral 5HT3 receptors (a type of serotonin receptor)
<p>Side effects</p>	 <p>Constipation Headache Dizzy</p>

	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Breastfeeding — OK for 1-2 doses</p> </div> <div style="text-align: center;">  <p>Children under 2 years</p> </div> <div style="text-align: center;">  <p>Liver trouble — use lower dose</p> </div> </div>
<p>Warnings</p>	<p>Interactions — Do not use together without medical consult</p> <p>ONDANSETRON with</p> <ul style="list-style-type: none"> • Apomorphine • Rifampicin • Tramadol — reduces analgesic effect of tramadol
<p>Tell the patient</p> 	<ul style="list-style-type: none"> • If too nauseous to swallow tablets or medicine mixture — dissolve wafer on the tongue
<p>Check</p>	<ul style="list-style-type: none"> • For cause of nausea