
















<p>Active ingredients (generic names)</p>	<p>PROBENECID (pro-ben-a-sid)</p> 
<p>What it is used for</p>	<p>Helps antibiotics work better in treatment of</p> <ul style="list-style-type: none"> • Skin infections <ul style="list-style-type: none"> ▸ Abnormal vaginal discharge ▸ Discharge from penis • Can use for long-term treatment of gout
<p>How it works</p>	<ul style="list-style-type: none"> • Reduces elimination of some acidic medicines (eg penicillins, cephalosporins) by the kidneys — so more antibiotic stays in body and works for a longer time • Increases removal of uric acid by the kidneys — reducing risk of gout attack
<p>Side effects</p>	<div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;">  <p>Rash</p> </div> <div style="text-align: center;">  <p>Nausea and vomiting</p> </div> <div style="text-align: center;">  <p>Headache</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;">  <p>Dizzy</p> </div> <div style="text-align: center;">  <p>Kidney stones</p> </div> </div>

 <p>Warnings</p>	<ul style="list-style-type: none"> • Do not start (or restart) use during a gout attack <ul style="list-style-type: none"> ▶ If person already taking regularly — can continue to use <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>Kidney stones</p> </div> <div style="text-align: center;">  <p>Pregnant</p> </div> </div> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>Breastfeeding</p> </div> <div style="text-align: center;">  <p>Children under 2 years</p> </div> </div>
	<p>Interactions — Do not use together without medical consult Probenecid interacts with a lot of different medicines — always check with doctor or pharmacist before adding another medicine</p>
<p>Tell the patient</p> 	<ul style="list-style-type: none"> • Do not take more than 1 aspirin a day unless doctor tells you to <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>Take with food or milk — so medicine doesn't make you feel sick</p> </div> <div style="text-align: center;">  <p>Drink lots of water — helps stop build up of uric acid, can prevent kidney stones</p> </div> </div> <p>Warning stickers (page 310): 10a, B</p>
<p>Check</p>	<div style="text-align: center;">  <p>Blood test Regular checks for kidney function (UEC), FBC</p> </div>